

Half-filled Kondo lattice on the honeycomb lattice

Yin Zhong,^{1,*} Ke Liu,² Yu-Feng Wang,¹ Yong-Qiang Wang,² and Hong-Gang Luo^{1,3,†}

¹*Center for Interdisciplinary Studies & Key Laboratory for Magnetism and Magnetic Materials of the MoE, Lanzhou University, Lanzhou 730000, China*

²*Institute of Theoretical Physics, Lanzhou University, Lanzhou 730000, China*

³*Beijing Computational Science Research Center, Beijing 100084, China*

(Dated: September 18, 2012)

The ground-state phase diagram of the Kondo lattice model on the honeycomb lattice at half-filling is carefully studied by using an extended mean-field theory. By treating magnetic interaction and Kondo screening on an equal footing, it is found that besides a trivial discontinuous first-order quantum phase transition between well-defined Kondo insulator and antiferromagnetic insulating state, there can exist a wide coexistent region with both Kondo screening and antiferromagnetic orders in the intermediate coupling regime. The appearance of such wide coexistence region is attributed to the unique linear density of state on the honeycomb lattice at half-filling in contrast to the case of square lattice. It is expected such a coexistence region could be realized by experiments of ultra-cold atoms on the honeycomb optical lattices and may be found by quantum Monte Carlo simulation in future since it is free of the ‘minus sign’. Additionally, the stability of Kondo insulator to the trivial decoupled state requires a minimum strength of the Kondo coupling, which is also the result of the magic linear density of state. We hope the present work may be helpful for further studies on the interplay between conduction electrons and the densely localized spins for the honeycomb lattice.

I. INTRODUCTION

It is still a challenge to understand the emergent quantum phases and corresponding quantum criticality in heavy fermion systems.^{1–12} To attack this challenging problem, the so-called Kondo lattice model is introduced, which is believed to capture the nature of interplay between Kondo screening and the magnetic interaction, namely, the Ruderman-Kittel-Kasuya-Yosida exchange interaction, mediated by conduction electrons among localized spins.¹³ The former effect favors a nonmagnetic spin singlet state in strong coupling limit while the latter tends to stabilize usual magnetic ordered states in weak coupling limit. There seems to exist a quantum phase transition or even a coexistence regime between these two kinds of well-defined states,^{14–19} however, a more radical critical phase may have been observed in $\text{YbRh}_2(\text{Si}_{0.95}\text{Ge}_{0.05})_2$.^{4,20}

The nature of these mentioned phenomena has been a long standing but controversial issue since the work of Doniach.²¹ It is noted that an extended mean-field theory treating the magnetic interaction and the Kondo screening on an equal footing, has predicted a coexistence regime of the disorder Kondo singlet and the antiferromagnetic ordered state with small staggered magnetization and partially screened local moments in the intermediate coupling.¹⁵ This coexistence has been confirmed by sophisticated quantum Monte Carlo (QMC) simulation without notorious ‘minus sign’ problem.^{16,17} Therefore, it is fair to say that such a mean-field theory could provide reliable physical results in the intermediate and strong Kondo coupling regimes when the unconventional quantum phases, e.g., quantum spin liquids, are irrelevant in the related problems.^{22,23}

Recently, much attention has been focused in strongly

correlated physics on the honeycomb lattice since the low energy excitations are described by relativistic Dirac fermions around distinct Dirac points rather than usual non-relativistic Landau quasiparticle near Fermi surface.^{24–35}

So far, most of the theoretical studies in this active field focus on Hubbard model or its derivative with spin-orbit coupling, the Kane-Mele-Hubbard.^{25,26,28,29,33–35} How the conduction electrons interplay with the densely localized spins on the honeycomb lattice, which could be encoded in terms of the Kondo lattice model in principle, is largely an open problem.³⁶

In this work, we try to uncover the ground-state phase diagram of the Kondo lattice model on the honeycomb lattice at half-filling using the mentioned extended mean-field theory for an anisotropic Kondo lattice model. Our main result is systematically summarized in Fig. 1. In weak Kondo coupling an antiferromagnetic ordered insulating state appears, and in strong coupling limit the Kondo insulator is ground state. In the intermediate regime, a wide coexistence regime is found. In some respect, our work may be considered as the first attempt in this direction before more sophisticated numerical approaches can be used.

II. THE KONDO LATTICE MODEL ON THE HONEYCOMB LATTICE AND MEAN-FIELD DECOUPLING

The model we considered is the anisotropic Kondo lattice model defined on the honeycomb lattice at half-




FIG. 1. A possible ground-state phase diagram of the Kondo lattice model on the honeycomb lattice at half-filling. AFM refers the antiferromagnetic ordered insulating state in weak Kondo coupling while Kondo insulator appears in strong coupling. In the intermediate regime, a coexistence region is found with both Kondo screening and antiferromagnetic orders.

filling,¹⁵

$$\begin{aligned}
 H &= H_t + H_{\parallel} + H_{\perp}, \\
 H_t &= -t \sum_{\langle ij \rangle \sigma} (c_{i\sigma}^{\dagger} c_{j\sigma} + c_{j\sigma}^{\dagger} c_{i\sigma}), \\
 H_{\parallel} &= \frac{J_{\parallel}}{4} \sum_i (c_{i\uparrow}^{\dagger} c_{i\uparrow} - c_{i\downarrow}^{\dagger} c_{i\downarrow}) (d_{i\uparrow}^{\dagger} d_{i\uparrow} - d_{i\downarrow}^{\dagger} d_{i\downarrow}), \\
 H_{\perp} &= \frac{J_{\perp}}{2} \sum_i (c_{i\uparrow}^{\dagger} c_{i\downarrow} d_{i\downarrow}^{\dagger} d_{i\uparrow} + c_{i\downarrow}^{\dagger} c_{i\uparrow} d_{i\uparrow}^{\dagger} d_{i\downarrow}), \quad (1)
 \end{aligned}$$

where H_t describes conduction electrons hopping between nearest-neighbor sites and the pseudofermion representation for local spins has been utilized as $S_i^{\alpha} = \frac{1}{2} \sum_{\sigma\sigma'} d_{i\sigma}^{\dagger} \tau_{\sigma\sigma'}^{\alpha} d_{i\sigma'}$ with τ^{α} being usual Pauli matrix and a local constraint $d_{i\uparrow}^{\dagger} d_{i\uparrow} + d_{i\downarrow}^{\dagger} d_{i\downarrow} = 1$ enforced in every site. $H_{\parallel}$ denotes the magnetic instability due to the polarization of conduction electrons by local spins while $H_{\perp}$ describes the local Kondo screening effect resulting from spin-flip scattering process of conduction electrons by local moments. The Kondo screening effect has also been formulated in the $1/N$ expansion but magnetic instability is difficult to treat in such a framework.³⁷

Now, using the mean-field decoupling introduced by Zhang and Yu¹⁵ for the longitudinal and transverse interaction term $H_{\parallel}$, $H_{\perp}$, respectively, it is straightforward to obtain a mean-field Hamiltonian

$$\begin{aligned}
 H_{MF} &= H_t + H_{\parallel}^{MF} + H_{\perp}^{MF} + E_0, \\
 H_t &= -t \sum_{k\sigma} (f(k) c_{kA\sigma}^{\dagger} c_{kB\sigma} + f^*(k) c_{kB\sigma}^{\dagger} c_{kA\sigma}), \\
 H_{\parallel}^{MF} &= \frac{J_{\parallel}}{2} \sum_{k\sigma} [\sigma (-m_c d_{kA\sigma}^{\dagger} d_{kA\sigma} + m_d c_{kA\sigma}^{\dagger} c_{kA\sigma}) \\
 &\quad - (A \rightarrow B)] \\
 H_{\perp}^{MF} &= \frac{J_{\perp} V}{2} \sum_{k\sigma} (c_{kA\sigma}^{\dagger} d_{kA\sigma} + c_{kB\sigma}^{\dagger} d_{kB\sigma} + h.c.), \\
 E_0 &= N_s (2J_{\parallel} m_d m_c + J_{\perp} V^2),
 \end{aligned}$$

where we have defined several mean-field parameters as $\langle d_{iA\uparrow}^{\dagger} d_{iA\uparrow} - d_{iA\downarrow}^{\dagger} d_{iA\downarrow} \rangle = 2m_d$, $\langle d_{iB\uparrow}^{\dagger} d_{iB\uparrow} - d_{iB\downarrow}^{\dagger} d_{iB\downarrow} \rangle = -2m_d$, $\langle c_{iA\uparrow}^{\dagger} c_{iA\uparrow} - c_{iA\downarrow}^{\dagger} c_{iA\downarrow} \rangle = -2m_c$, $\langle c_{iB\uparrow}^{\dagger} c_{iB\uparrow} -$

$c_{iB\downarrow}^{\dagger} c_{iB\downarrow} \rangle = 2m_c$ and $-V = \langle c_{i\uparrow}^{\dagger} d_{i\uparrow} + d_{i\downarrow}^{\dagger} c_{i\downarrow} \rangle = \langle c_{i\downarrow}^{\dagger} d_{i\downarrow} + d_{i\uparrow}^{\dagger} c_{i\uparrow} \rangle$ with $f(k) = e^{-ik_x} + 2e^{ik_x/2} \cos \sqrt{3}k_y/2$ and A , B representing two nonequivalent sublattices. It can be seen that m_d, m_c corresponds to magnetization of local spins and conduction electrons, respectively, while non-vanishing V denotes the onset of Kondo screening effect. Besides, since we are considering a half-filled lattice, the local constraint has been safely neglected at the present mean-field level with chemical potential setting to zero.¹⁵

Diagonalizing the above mean-field Hamiltonian, the four quasiparticle bands are obtained as

$$E_{\pm\pm}(k) = \pm \frac{1}{\sqrt{2}} \sqrt{F_1(k) \pm F_2(k)} \quad (2)$$

with $F_1(k) = J_{\parallel}^2(m_d^2 + m_c^2)/4 + J_{\perp}^2 V^2/2 + t^2 |f(k)|^2$ and $F_2(k) = [F_1(k)^2 - 4(J_{\parallel}^4 m_d^2 m_c^2/16 + J_{\parallel}^2 m_d m_c J_{\perp}^2 V^2/8 + J_{\perp}^4 V^4/16) - J_{\parallel}^2 m_c^2 t^2 |f(k)|^2]^{\frac{1}{2}}$. Therefore, the wanted ground-state energy at half-filling is given by

$$E_g = \sum_k (E_{--}(k) + E_{-+}(k)) + E_0. \quad (3)$$

III. THE MEAN-FIELD SOLUTION FOR GROUND-STATE

Firstly, we proceed to discuss two simple but physically interesting limits for Kondo coupling $J_{\parallel}$ and $J_{\perp}$, which correspond to the antiferromagnetic ordered state ($J_{\parallel} \gg J_{\perp}$) and Kondo insulating state ($J_{\parallel} \ll J_{\perp}$), respectively.¹³

For the case with $J_{\parallel} \gg J_{\perp}$, in general, one expects the antiferromagnetic ordered state to be the stable ground-state of Kondo lattice model on the honeycomb lattice due to its bipartite feature.¹³ To study the possible antiferromagnetic ordered state, making use of the ground-state energy (Eq.3) with assuming no Kondo screening existing ($V = 0$) and the quasiparticle spectrum (Eq.2), we can easily derive ground-state energy of the antiferromagnetic ordered state per site as

$$E_g^{AFM} = J_{\parallel} m_c (2m_d - 1) - \frac{1}{N_s} \sum_k \sqrt{J_{\parallel}^2 m_d^2 + 4t^2 |f(k)|^2}.$$

and two self-consistent equations from minimizing E_g^{AFM} with respect to magnetization m_d and m_c , respectively.

$$\begin{aligned}
 J_{\parallel} m_c (2m_d - 1) &= 0, \\
 2J_{\parallel} m_c - \frac{J_{\parallel}^2 m_d}{2\Lambda^2} [(4\Lambda^2 + J_{\parallel}^2 m_d^2) - J_{\parallel} m_d] &= 0
 \end{aligned}$$

where we have used a simplified linear density of state (DOS) $\rho(\varepsilon) = |\varepsilon|/\Lambda^2$ when transforming the summation over momentum k into integral on energy ε with Λ being high-energy cutoff.²⁴ Thus, $t|f(k)|$ can be replaced by $|\varepsilon|$ to simplify corresponding calculations.

From these two equations, one obtains $m_d = 1/2$ and $m_c = J_{\parallel} [(4\Lambda^2 + J_{\parallel}^2/4) - J_{\parallel}/2]/8$. Meanwhile,

the low-lying quasiparticle excitations in the antiferromagnetic ordered state has the energy $E_{\pm+}(k) = \pm\sqrt{t^2|f(k)|^2 + J_{\parallel}^2/16}$ and $E_{\pm-}(k) = \pm J_{\parallel}m_c/2$ with an apparent gap $J_{\parallel}/4$ around the Dirac points where $f(k) = 0$. Thus, we conclude the antiferromagnetic ordered state we obtained is indeed an insulating state with fully polarized local spins ($m_d = 1/2$) while conduction electrons only partially polarize ($m_c < 1/2$). This feature is similar to the previous study on square lattice, thus confirms the validity of our current treatment.¹⁵

Another interesting case appears when $J_{\parallel} \ll J_{\perp}$. It is natural to expect that a Kondo insulating state arises in this situation for half-filling.^{13,15} Following the same methodology of treating antiferromagnetic insulating state, we can get the ground-state energy per site for the expected Kondo insulating state with $V \neq 0$ but no magnetic orders $m_d = m_c = 0$

$$E_g^{Kondo} = J_{\perp}V^2 - \frac{4}{3\Lambda^2}[(\Lambda^2 + J_{\perp}^2V^2)^{3/2} - J_{\perp}^3V^3].$$

Minimizing E_g^{Kondo} with respect to Kondo hybridization parameter V , we obtain $V = 1 - \Lambda^2/(4J_{\perp}^2)$ which implies a critical coupling $J_{\perp}^c = 1/2$ corresponding to vanishing V . It is noted that $V \propto (J_{\perp} - J_{\perp}^c)$ in contrast to usual mean-field result $\beta = 1/2$, and this can be attributed to the low-energy linear DOS of conduction electrons on the honeycomb lattice at half-filling. Similar critical behavior for onset of Kondo screening on the honeycomb lattice has been obtained in the study of Kondo breakdown mechanism as well.²³ As a matter of fact, the existence of the critical coupling $J_{\perp}^c = 1/2$ results from the competition between the Kondo insulating state and the trivial decoupled state where $V = m_d = m_c = 0$, its ground-state energy $E_g^0 = -4\Lambda/3$ comes solely from free conduction electrons. Comparing E_g^0 and E_g^{Kondo} , one clearly recovers the critical coupling $J_{\perp}^c$, which justifies the above simple picture.

However, since the Kondo insulating state is unstable to the decoupled state when $J_{\perp} < J_{\perp}^c = 1/2$, one may wonder whether a trivial decoupled state appears between the Kondo insulating state and the antiferromagnetic insulating state. It is easy to see the ground-state energy of the antiferromagnetic state E_g^{AFM} is always lower than E_g^0 for any positive coupling $J_{\parallel}$. Therefore, a trivial disordered state in intermediate coupling seems unfavorable based on our current mean-field treatment.

Instead, we may consider a possible quantum critical point (QCP) between Kondo insulating state and the antiferromagnetic state and its position can be determined by comparing the ground-state energies of E_g^{AFM} and E_g^{Kondo} . For physically interesting case with $J_{\perp} = J_{\parallel} = J$, we have

$$\begin{aligned} \frac{1}{6\Lambda^2}[(4\Lambda^2 + J^2/4)^{\frac{3}{2}} - (J^2/4)^{\frac{3}{2}}] &= \frac{-1}{J}(J - \frac{\Lambda^2}{4J^2})^2 \\ + \frac{4}{3\Lambda^2}[(\Lambda^2 + (J - \frac{\Lambda^2}{4J^2})^2)^{\frac{3}{2}} - (J - \frac{\Lambda^2}{4J^2})^3]. \end{aligned} \quad (4)$$

The position of the QCP is readily obtained as $J_c = 0.88\Lambda$, numerically. However, in fact, one can check that this putative QCP is a first-order quantum phase transition point when comparing the first-order derivative of E_g^{Kondo} and E_g^{AFM} with respect to the Kondo coupling J . Thus, we do not expect radical critical behaviors near such first-order quantum phase transition point in the spirit of Landau-Ginzburg-Wilson paradigm.¹ Additionally, it is noted that such first-order quantum phase transition has also been obtained on the square lattice³⁸ and we suspect this feature may be generic for conventional mean-field treatment of Kondo lattice models according to standard Landau-Ginzburg phase transition theory.¹⁵ However, we point out that it is a subtle issue to compare the results of a first-order quantum transition with numerical simulations, particularly when the first-order transition is a weak one.³⁹

Generically, in the intermediate coupling regime, a possible coexistence region of the Kondo insulating state and the antiferromagnetic ordered state cannot be excluded.¹⁵ For searching such possibility, we have to resort to the full formalism of ground-state energy E_g (Eq.3) and three self-consistent equations derived from $\frac{\partial E_g}{\partial V} = \frac{\partial E_g}{\partial m_d} = \frac{\partial E_g}{\partial m_c} = 0$. Since the possible coexistence region has two boundaries, which correspond to the onset of Kondo screening ($V \neq 0$) in antiferromagnetic insulating state and antiferromagnetic order ($m_c, m_d \neq 0$) arising in Kondo insulator, we can derive two equations for these two distinct boundaries.

The first boundary which means the antiferromagnetic order ($m_c, m_d \neq 0$) arising in Kondo insulator can be obtained by solving the following equation

$$\begin{aligned} J(1 - \frac{\Lambda^2}{4J^2})^2 &= \frac{2}{3\Lambda^2}(\Lambda^2 - J^2 - \frac{\Lambda^4}{16J^2})\sqrt{\Lambda^2 + J^2(1 - \frac{\Lambda^2}{4J^2})^2} \\ + \frac{2}{3\Lambda^2}J^3(1 - \frac{\Lambda^2}{4J^2})^3, \end{aligned} \quad (5)$$

where we have utilized the expression for the Kondo hybridization parameter $V = 1 - \Lambda^2/(4J_{\perp}^2)$ and assuming the isotropic condition ($J_{\perp} = J_{\parallel} = J$). Solving Eq.5 numerically, one finds the first boundary located in $J_{c1} = 0.78\Lambda$ which is smaller than first-order quantum phase transition point ($J_c = 0.88\Lambda$), thus the expected coexistent scenario is stable to the previous discontinuous first-order quantum phase transition.

For the second boundary, which corresponds to the onset of Kondo screening ($V \neq 0$) in antiferromagnetic state, one can also derive similar equation like Eq.(5) to determine the location of the boundary.

$$\frac{2\Lambda^2}{J} = \int_0^{\Lambda} d\varepsilon [P(\varepsilon) + Q(\varepsilon) + (P(\varepsilon) - Q(\varepsilon))W(\varepsilon)], \quad (6)$$

where we have defined three auxiliary functions $P(\varepsilon) = 1/\sqrt{\varepsilon^2 + J^2/16}$, $Q(\varepsilon) = 2/Jm_c$ and $W(\varepsilon) = [\varepsilon^2 + J^2/16 + J^2(m_c^2 - m_c)/4]/[\varepsilon^2 + J^2(1/4 - m_c^2)/4]$ with $m_c = J_{\parallel}[(4\Lambda^2 + J_{\parallel}^2/4) - J_{\parallel}/2]/8$. Then, one obtains the location

of the second boundary $J_{c2} = 1.06\Lambda$ by numerically solving Eq.(6). It is noted that $J_{c2} = 1.06\Lambda > J_c = 0.88\Lambda$, which implies the system indeed tends to form a coexistent state with Kondo screening and antiferromagnetic orders.

Therefore, we may conclude that there may exist a coexistence regime ($J_{c1} = 0.78\Lambda < J < J_{c2} = 1.06\Lambda$) in the intermediate coupling regime based on the discussion in two phase boundaries for the isotropic case ($J_{\perp} = J_{\parallel} = J$). When comparing to the square lattice, we find the range of the coexistence regime on the honeycomb lattice is obviously larger than the one on the square lattice with $J_{c1} = 0.56\Lambda < J < J_{c2} = 0.62\Lambda$.¹⁵ We suspect this difference may result from unique linear DOS on the honeycomb lattice at half-filling, which lowers the energy penalty of coexistence.

IV. EXTENSIONS AND DISCUSSIONS

Moreover, for physical observable, it is expected that in the proposed coexistence regime the dynamical magnetic structure factor, which can be detected in inelastic neutron scattering experiments, should have two contributions. One is from a momentum-independent single site slow component due to the localized Kondo-type excitations, and the other is contributed from a strongly momentum-dependent inter-site fast component, reflecting the magnetic interactions.¹⁵

We noted that, it is also interesting to study a more simplified model only including degrees of freedom for spins, namely, the Kondo necklace model on the honeycomb lattice.²¹ However, the present mean-field decoupling approach is not cheap in such a spin model but the so-called bond-operator representation could be a useful tool in this case.⁴⁰ Moreover, if a spin-orbit coupling is introduced in our present model as the case in Ref. [36], one could study the interplay among the antifer-

romagnetic order, Kondo screening and the symmetry-protected topological states (such as the 2D topological insulator). But, it seems that the topological insulating state found in Ref. [36] might be unstable to the conventional antiferromagnetic order in terms of the present mean-field decoupling treatment.⁴¹ Therefore, more sophisticated treatments on this issue is eagerly desirable.

V. CONCLUSION

In summary, we have obtained the ground-state phase diagram of the Kondo lattice model on the honeycomb lattice at half-filling using extended mean-field decoupling. By treating magnetic interaction and Kondo screening on an equal footing, it is found that besides a trivial discontinuous first-order quantum phase transition between well-defined Kondo insulator and antiferromagnetic insulating state, there can exist a wide coexistence regime with both Kondo screening and antiferromagnetic orders in the intermediate coupling regime. It is expected such a coexistence regime could be realized by experiments of ultra-cold atoms on the honeycomb optical lattices and may be found by quantum Monte Carlo simulation in future since it is free of the ‘minus sign’.^{16,17,42,43} We hope the present work may be helpful for further studies on the interplay between conduction electrons and the densely localized spins for the honeycomb lattice.

ACKNOWLEDGMENTS

The work was supported partly by NSFC, the Program for NCET, the Fundamental Research Funds for the Central Universities and the national program for basic research of China.

* zhongy05@hotmail.com

† luohg@lzu.edu.cn

¹ S. Sachdev, *Quantum Phase Transition*, 2nd ed. (Cambridge University Press, Cambridge, England, 2011).

² H. V. Löhneysen, A. Rosch, M. Vojta and P. Wölfle, *Rev. Mod. Phys.* **79**, 1015 (2007).

³ M. Vojta, *J Low Temp Phys* **161**, 203 (2010).

⁴ J. Custers et al., *Nature (London)* **424**, 524 (2003).

⁵ J. Custers, P. Gegenwart, C. Geibel, F. Steglich, P. Coleman and S. Paschen, *Phys. Rev. Lett.* **104**, 186402 (2010).

⁶ Y. Matsumoto et al., *Science* **331**, 316 (2011).

⁷ T. Senthil, S. Sachdev and M. Vojta, *Phys. Rev. Lett.* **90**, 216403 (2003).

⁸ T. Senthil, M. Vojta and S. Sachdev, *Phys. Rev. B* **69**, 035111 (2004).

⁹ C. Pépin, *Phys. Rev. Lett.* **94**, 066402 (2005).

¹⁰ K. S. Kim and C. L. Jia, *Phys. Rev. Lett.* **104**, 156403 (2010).

¹¹ T. Grover and T. Senthil, *Phys. Rev. B* **81**, 205102 (2010).

¹² Y. Zhong, K. Liu, Y. Q. Wang and H.-G. Luo, *Phys. Rev. B* **86**, 115113 (2012).

¹³ H. Tsunetsugu, M. Sigrist and K. Ueda, *Rev. Mod. Phys.* **69**, 809 (1997).

¹⁴ C. Lacroix and M. Cyrot, *Phys. Rev. B* **20**, 1969 (1979).

¹⁵ G. M. Zhang and L. Yu, *Phys. Rev. B* **62**, 76 (2000).

¹⁶ S. Capponi and F. F. Assaad, *Phys. Rev. B* **63**, 155114 (2001).

¹⁷ H. Watanabe and M. Ogata, *Phys. Rev. Lett.* **99**, 136401 (2007).

¹⁸ G.-B. Li and G.-M. Zhang, *Phys. Rev. B* **81**, 094420 (2010).

¹⁹ G.-M. Zhang, Y.-H. Su, and Lu Yu, *Phys. Rev. B* **83**, 033102 (2011).

²⁰ J. Custers, P. Gegenwart, C. Geibel, F. Steglich, P. Coleman and S. Paschen, *Phys. Rev. Lett.* **104**, 186402 (2010).

²¹ S. Doniach, *Physica B and C* **91**, 231 (1977).

- ²² P. A. Lee, N. Nagaosa and X. G. Wen, Rev. Mod. Phys. **78**, 17 (2006).
- ²³ S. Saremi and P. A. Lee, Phys. Rev. B **75**, 165110 (2007).
- ²⁴ A. H. C. Neto, F. Guinea, N. M. R. Peres, K. S. Novoselov and A. K. Geim, Rev. Mod. Phys. **81**, 109 (2009).
- ²⁵ V. N. Kotov, B. Uchoa, V. M. Pereira, F. Guinea and A. H. C. Neto, arXiv:cond-mat/1012.3484 (2010). (To publish in Rev. Mod. Phys.)
- ²⁶ Z. Y. Meng, T. C. Lang, S. Wessel, F. F. Assaad, and A. Muramatsu, Nature (London) **464**, 847 (2010).
- ²⁷ I. F. Herbut, Phys. Rev. Lett. **97**, 146401 (2006).
- ²⁸ B. K. Clark, D. A. Abanin and S. L. Sondhi, Phys. Rev. Lett. **107**, 087204 (2011).
- ²⁹ F. Mezzacapo and M. Boninsegni, Phys. Rev. B **85**, 060402(R) (2012).
- ³⁰ Y. Zhong, K. Liu, Y. Q. Wang and H.-G. Luo, arXiv:cond-mat/1205.6064 (2012).
- ³¹ M. Z. Hasan and C. L. Kane, Rev. Mod. Phys. **82**, 3045 (2010).
- ³² X.-L. Qi and S.-C. Zhang, Rev. Mod. Phys. **83**, 1057 (2011).
- ³³ S. Rachel and K. Le Hur, Phys. Rev. B **82**, 075106 (2010).
- ³⁴ M. Hohenadler, T.-C. Lang and F. F. Assaad, Phys. Rev. Lett. **106** 100403 (2011).
- ³⁵ A. Rüegg and G. A. Fiete, Phys. Rev. Lett. **108** 046401 (2012).
- ³⁶ X.-Y. Feng, J. Dai, C.-H. Chung and Qimiao Si, arXiv:cond-mat/1206.0979v1 (2012).
- ³⁷ S. Doniach, Phys. Rev. B **35**, 1814 (1987).
- ³⁸ Using the ground-state energy of antiferromagnetic state (Eq.7) and Kondo insulator (Eq.8) in Ref. [15], it is easy to check that at the putative QCP $J_c = 0.58D$, there exists a jump about 0.1 for the first-order derivative of ground-state energy of antiferromagnetic state and Kondo insulator with respect to the Kondo coupling. Thus, a first-order transition is clearly observed in the mean-field treatment of Ref. [15].
- ³⁹ M. A. Continentino, *Quantum Scaling in Many-Body Systems* .(World Scientific Press, Singapore, 2001).
- ⁴⁰ G.-M. Zhang, Q. Gu and L. Yu, Phys. Rev. B **62**, 69 (2000).
- ⁴¹ Y. Zhong, Y.-F. Wang, Y.-Q. Wang and H.-G. Luo, in preparing. We have studied the model (Eq.1) in Ref. [36] in terms of the mean-field decoupling described in the main text. We found the topological insulating state ($V = m_c = m_d = 0$) always has higher ground-state energy than the usual antiferromagnetic state ($V = 0, m_c = m_d \neq 0$) in spite of the fact that the topological insulating state is more stable than the trivial Kondo insulating state in weak Kondo coupling regime. Therefore, based on our mean-field results, the quantum phase transition studied in Ref. [36] might give way to a trivial first-order transition between the antiferromagnetic state and Kondo insulating state if no other kinds of interaction or hopping terms are added.
- ⁴² I. Bloch, J. Dalibard and W. Zwerger, Rev. Mod. Phys. **80**, 885 (2008).
- ⁴³ N. Goldman, A. Kubasiak, A. Bermudez, P. Gaspard, M. Lewenstein, and M. A. Martin-Delgado, Phys. Rev. Lett. **103**, 035301 (2009).